



Personal Data

Name : BG_54
Sex : female
Age : 54

Numerical Data Analysis

PARAMETERS OF THE IRIS :	S	D
Diameter (pix) =	416	424
Area (pix)	135918	141196
PARAMETERS OF THE PUPILLARY BORDER :		
Diameter (pix)	102	96
Diameter of the pupil in relation to iris (%)	24	22
Normal for current age 21-25%	Normal	Normal
Pupil border deformation degree (normal:0%...5%)	1	7
PARAMETERS OF THE PUPIL RELATIVE TO THE IRIS :		
Distance between the pupil and iris centers (%)	4.81	4.25
Normal (lower than 5% of above) or pathology	Normal	Normal
PARAMETERS OF THE APPROXIMATE ELLIPSE		
Ellipseness degree of the pupil (normal: 95% ... 100%)	91	98
Pupil form type	Pathology ellipse	Normal circle
PARAMETERS OF THE PUPILLARY MARGIN :		
Type of the form -	regular Normal	regular Normal

Diagnosis

S : Frontal drawing in (11:36 - 0:46) - 1.96 %
S : Middle-temporal protrusion (2:02 - 3:56) - 3.92 %
S : Middle-nasal protrusion (7:46 - 9:24) - 1.96 %
S : Decentralization of the pupil is normal.
S : Oval-horizontal form of the pupil.

D : Frontal flatness (10:50 - 0:26) - 4.17 %
D : Decentralization of the pupil is normal.
D : Ellipseness of the pupil is normal.

Discirculatory encephalopathy. Depression.
Depression, blood cerebral circulation disturbance, Bronchial asthma
predisposition.

PARAMETERS OF THE AUTONOMIC NERVE WREATH (ANW):	S	D
Diameter (pix)	212	212
Perimeter (pix)	696	733
The ratio between Pupillary and Ciliary belts (%)	35.03	35.37
Normal (25..35%) or pathologic.	Atonic	Atonic
Asymmetry of pupillary belt (normal: 0..5%)	4.72	6.60
	Normal	Pathology
Type of the ANW form -	regular	drawn in
	Normal	Pathology

D: Lower temporal shift.
D: Frontal zone of pupillary belt is constricted.
S: Frontal zone of pupillary belt is constricted.

Sympathotonic. Liminal sensitivity of nervous system is decreased. Reflex
activity is decelerated. Secretory and evacuation functions of digestive
tract is reduced.
Hemodynamics disturbance in the vena cava inferior.
Nephroptosis. (stable changes).

Vertebral osteoarthrosis.

HEMATOGENIC SUBTYPE.

Decreased reactivity of lymphatic system, leukocytopenia in infectious diseases, microelement deficiency. Predisposition to lymphadenitis and adenomas.

Metabolic disturbances : osteochondrosis, arthrosis, lithogenesis.

ADVICE

Preventive measures should be directed to increase of lymphatic system reactivity and prevention of metabolic violations.

1. Limitation of products, reach with topo-stable calcium.
2. Strengthening of the organism.
3. Prevention of virus hepatitis.
4. Regular, subdivided nutrition.
5. Great care should be given to personal hygiene, because of predisposition to infections diseases.
6. Active regular gymnastics, directed the strengthening of back muscles.
7. Fruit-vegetable cocktails, rich with PP and B group vitamins.
8. Introductory nature therapy (it should be carried out under doctor's control) :

PURIFICATION OF JOINTS : 5 g of laurel leaves are put into 300 g of water and boil within 5 min. After it the liquid should be brewed in a thermos for 3-4 hours. Take the solution by sips with the intervals within 12 hours (never drink it at once - haemorrhaging may be result!).

Continue this procedure for 3 days. You may repeat it in a week.

Purification of joints can be carried out once every 3 months the first year, then once a

year. This procedure will help you to be free from salt deposits in the joints, from "weather" pains, joint fatigue, etc.

9. Phytotherapy:

Tea

1 table spoon of oats (30 g) pour with 300 ml of cold boiled water, soak within 2 hours. On the lower fire drive to boiling in the same water, boil within 20 min, filter into the thermos

1 tea spoon of the dill seeds pour with 100 ml of boiling water and put on the water bath for 15 min. After that filter into the same thermos.

All preparations should be done in the evening. In the morning and in the afternoon take 200 ml of the solution 10-15 min before meals. The course consists of 10-12 days, it should be repeated twice a years.