



Personal Data

Name : BR51  
Sex : male  
Age : 51

Numerical Data Analysis

PARAMETERS OF THE IRIS :	S	D
Diameter (pix) =	420	436
Area (pix)	138544	149301
PARAMETERS OF THE PUPILLARY BORDER :		
Diameter (pix)	108	106
Diameter of the pupil in relation to iris (%)	25	24
Normal for current age 21-25%	Normal	Normal
Pupil border deformation degree (normal:0%...5%)	3	0
PARAMETERS OF THE PUPIL RELATIVE TO THE IRIS :		
Distance between the pupil and iris centers (%)	7.14	2.75
Normal (lower than 5% of above) or pathology	Pathology	Normal
PARAMETERS OF THE APPROXIMATE ELLIPSE		
Ellipseness degree of the pupil (normal: 95% ... 100%)	93	92
Pupil form type	Pathology ellipse	Pathology ellipse
PARAMETERS OF THE PUPILLARY MARGIN :		
Type of the form -	regular Normal	regular Normal

S : Upper nasal flatness ( 8:42 - 1:00) - 5.56 %  
 S : Middle-temporal protrusion ( 2:42 - 3:54) - 1.85 %  
 S : Middle-nasal decentralization.  
 S : Oval-horizontal form of the pupil.

D : Upper temporal flatness ( 9:08 - 0:32) - 3.77 %  
 D : Lower temporal protrusion ( 7:04 - 8:28) - 1.89 %  
 D : Decentralization of the pupil is normal.  
 D : Oval-diagonal form of the pupil.

The injury of cervical part of spinal column (etiology should be specified)  
 Risk to cardiospastic reaction.  
 Urogenital disturbances.  
 Depression, blood cerebral circulation disturbance, Bronchial asthma  
 predisposition.

PARAMETERS OF THE AUTONOMIC NERVE WREATH (ANW):	S	D
Diameter (pix)	194	188
Perimeter (pix)	804	698
The ratio between Pupillary and Ciliary belts (%)	27.56	24.85
Normal (25..35%) or pathologic.	Normal	Spastic
Asymmetry of pupillary belt (normal: 0..5%)	7.22	7.45
Type of the ANW form -	Pathology	Pathology
	drawn in lacerated	
	Pathology	Pathology

D: Lower temporal shift.  
 D: Frontal and basal zones of pupillary belt are constricted.  
 S: Lower temporal shift.  
 S: Frontal zone of pupillary belt is constricted.

Functional frustration in correlation between sympathetic and parasympathetic nervous systems. (The origin should be specified?)  
 Hemodynamics disturbance in the vena cava inferior.  
 Acquired colitis. Initial manifestations of pathology in compensation state.  
 Nephroptosis. (stable changes).

Changes in liver stroma. Decrease of detoxicative function.  
 Predisposition to allergic reactions.  
 Initial vertebral osteoarthritis.

LYMPHATIC HYDROGENOUS SUBTYPE WITH SIGNS OF URATE DIATHESIS.  
 Violation of colloid balance in protein liquids, produced by the organism.  
 Lymphoid tissue is highly reactive. Diseases, connected with metabolic disturbance are characteristic for the given subtype.  
 Diseases, connected with metabolic disturbances, are typical: arthrosis, spinal osteochondrosis, urolithic and cholelithic diseases, podagra.