



Personal Data

Name : Darryl
Sex : male
Age : 32

Numerical Data Analysis

PARAMETERS OF THE IRIS :	S	D
Diameter (pix) =	474	448
Area (pix)	176460	157633
PARAMETERS OF THE PUPILLARY BORDER :		
Diameter (pix)	134	132
Diameter of the pupil in relation to iris (%)	28	29
Normal for current age 23-32%	Normal	Normal
Pupil border deformation degree (normal:0%...5%)	9	11
PARAMETERS OF THE PUPIL RELATIVE TO THE IRIS :		
Distance between the pupil and iris centers (%)	2.95	6.70
Normal (lower than 5% of above) or pathology	Normal	Pathology
PARAMETERS OF THE APPROXIMATE ELLIPSE		
Ellipseness degree of the pupil (normal: 95% ... 100%)	92	97
Pupil form type	Pathology ellipse	Normal chord
	Pathology	Pathology
PARAMETERS OF THE PUPILLARY MARGIN :		
Type of the form -	regular	regular
	Normal	Normal

Diagnosis

S : Basal protrusion (4:34 - 6:48) - 2.99 %
S : Middle-nasal flatness (8:14 - 10:32) - 5.97 %
S : Decentralization of the pupil is normal.
S : Oval-vertical form of the pupil.

D : Basal protrusion (3:54 - 7:14) - 3.03 %
D : Upper-nasal decentralization.
D : Ellipseness of the pupil is normal.

Increased fatiguability.

Liver disease.

Circulatory cerebral disturbance with risk of spastic variation.

PARAMETERS OF THE AUTONOMIC NERVE WREATH (ANW):	S	D
Diameter (pix)	248	244
Perimeter (pix)	835	847
The ratio between Pupillary and Ciliary belts (%)	33.53	35.44
Normal (25..35%) or pathologic.	Normal	Atonic
Asymmetry of pupillary belt (normal: 0..5%)	13.71	4.92
	Pathology	Normal
Type of the ANW form -	drawn in lacerated	Pathology Pathology

D: Frontal zone of pupillary belt is constricted.

S: Lower temporal shift.

S: Frontal zone of pupillary belt is constricted.

Functional frustration in correlation between sympathetic and parasympathetic nervous systems. (The origin should be specified?)

Weakness of smooth muscles. (stable changes).

Changes in liver stroma. Decrease of detoxicative function.

Diffuse changes in pancreas.

Endogenic or exogenic intoxication.

Initial vertebral osteoarthrosis.

LYMPHATIC SUBTYPE.

Increased reactivity of lymphoid tissues (thymus, lymphatic nodes, tonsils, spleen, etc).

Predisposition to gastrointestinal diseases, diseases of genitals and those connected with immune system pathology (allergic diseases, collagenosis, reduced resistance to infections).