



Personal Data

Name : 453672  
 Sex : male  
 Age : 50

Numerical Data Analysis

	S	D
PARAMETERS OF THE IRIS :		
Diameter (pix) =	424	376
Area (pix)	141196	111036
PARAMETERS OF THE PUPILLARY BORDER :		
Diameter (pix)	82	84
Diameter of the pupil in relation to iris (%)	19	22
Normal for current age 21-25%	Miosis	Normal
Pupil border deformation degree (normal:0%...5%)	22	12
PARAMETERS OF THE PUPIL RELATIVE TO THE IRIS :		
Distance between the pupil and iris centers (%)	3.30	3.19
Normal (lower than 5% of above) or pathology	Normal	Normal
PARAMETERS OF THE APPROXIMATE ELLIPSE		
Ellipseness degree of the pupil (normal: 95% ... 100%)	95	87
Pupil form type	Normal chord	Pathology ellipse
	Pathology	Pathology
PARAMETERS OF THE PUPILLARY MARGIN :		
Type of the form -	regular	regular
	Normal	Normal

S : Decentralization of the pupil is normal.

S : Ellipseness of the pupil is normal.

D : Frontal drawing in (11:18 - 0:50) - 2.38 %

D : Middle-temporal protrusion ( 7:46 - 9:24) - 4.76 %

D : Decentralization of the pupil is normal.

D : Oval-horizontal form of the pupil.

Lungs disease in history.

Depression, blood cerebral circulation disturbance, Bronchial asthma  
predisposition.

PARAMETERS OF THE AUTONOMIC NERVE WREATH (ANW):	S	D
Diameter (pix)	202	162
Perimeter (pix)	677	576
The ratio between Pupillary and Ciliary belts (%)	35.09	26.71
Normal (25..35%) or pathologic.	Atonic	Normal
Asymmetry of pupillary belt (normal: 0..5%)	5.94	2.47
Type of the ANW form -	Pathology indented	Normal drawn in Pathology

D: Frontal zone of pupillary belt is constricted.

S: Middle-nasal shift.

S: Frontal and basal zones of pupillary belt are constricted.

Functional frustration in correlation between sympathetic and parasympathetic nervous systems. (The origin should be specified?)

Hypofunction of n. vagus or stellate ganglion.

Acquired colitis. Initial manifestations of pathology in compensation state.

Nephroptosis. (stable changes).

Vertebral osteoarthritis.

Metabolism is disturbed. Intoxication.