



Personal Data

Name : AM_36
 Sex : male
 Age : 36

Numerical Data Analysis

PARAMETERS OF THE IRIS :	S	D
Diameter (pix) =	474	462
Area (pix)	176460	167639
PARAMETERS OF THE PUPILLARY BORDER :		
Diameter (pix)	116	114
Diameter of the pupil in relation to iris (%)	24	24
Normal for current age 23-32%	Normal	Normal
Pupil border deformation degree (normal:0%...5%)	2	2
PARAMETERS OF THE PUPIL RELATIVE TO THE IRIS :		
Distance between the pupil and iris centers (%)	4.64	4.33
Normal (lower than 5% of above) or pathology	Normal	Normal
PARAMETERS OF THE APPROXIMATE ELLIPSE		
Ellipseness degree of the pupil (normal: 95% ... 100%)	93	95
Pupil form type	Pathology ellipse Pathology	Normal ellipse Pathology
PARAMETERS OF THE PUPILLARY MARGIN :		
Type of the form -	regular Normal	regular Normal

S : Frontal flatness (10:40 - 1:28) - 3.45 %
 S : Middle-temporal protrusion (2:28 - 3:48) - 3.45 %
 S : Middle-nasal flatness (7:52 - 9:28) - 15.52 %
 S : Decentralization of the pupil is normal.
 S : Oval-diagonal form of the pupil.

D : Frontal flatness (11:08 - 0:54) - 3.51 %
 D : Middle-temporal flatness (8:36 - 10:28) - 8.77 %
 D : Decentralization of the pupil is normal.
 D : Ellipseness of the pupil is normal.

Tonsillo-cardial syndrome.
 Increased fatiguability.
 Urogenital disturbances.

PARAMETERS OF THE AUTONOMIC NERVE WREATH (ANW):	S	D
Diameter (pix)	238	236
Perimeter (pix)	836	815
The ratio between Pupillary and Ciliary belts (%)	34.08	35.06
Normal (25..35%) or pathologic.	Normal	Atonic
Asymmetry of pupillary belt (normal: 0..5%)	5.88	0.85
Type of the ANW form -	Pathology	Normal
	drawn in	indented
	Pathology	Pathology

D: Frontal and basal zones of pupillary belt are constricted.
 S: Lower nasal shift.
 S: Frontal zone of pupillary belt is constricted.

Functional frustration in correlation between sympathetic and parasympathetic nervous systems. (The origin should be specified?)
 Atonic colitis. Initial manifestations of pathology in compensation state.
 Weakness of smooth muscles. (stable changes).

Increased emotional lability, predisposition to spastic reactions. Overstressed state.

PIGMENT SATURATED SUBTYPE.

For people with such subtype increased reactivity of lymphoid tissues (thymus, lymphatic nodes, tonsils, spleen, etc.) is typical.
 Diseases of digestive tract mucosa, genitals are often observed.
 Predisposition to immune system pathology (allergic diseases, collagenosis, reduced resistance to infections). Fast acetilator - sensitivity of the ferment systems to chemical and drug influences is increased.