



Personal Data

Name : AYC40
 Sex : female
 Age : 40

Numerical Data Analysis

PARAMETERS OF THE IRIS :	S	D
Diameter (pix) =	442	448
Area (pix)	153439	157633
PARAMETERS OF THE PUPILLARY BORDER :		
Diameter (pix)	102	102
Diameter of the pupil in relation to iris (%)	23	22
Normal for current age 23-32%	Normal	Miosis
Pupil border deformation degree (normal:0%...5%)	2	6
PARAMETERS OF THE PUPIL RELATIVE TO THE IRIS :		
Distance between the pupil and iris centers (%)	3.62	4.02
Normal (lower than 5% of above) or pathology	Normal	Normal
PARAMETERS OF THE APPROXIMATE ELLIPSE		
Ellipseness degree of the pupil (normal: 95% ... 100%)	98	96
	Normal	Normal
Pupil form type	circle	chord
	Normal	Pathology
PARAMETERS OF THE PUPILLARY MARGIN :		
Type of the form -	regular	regular
	Normal	Norma

Diagnosis

S : Decentralization of the pupil is normal.

S : Ellipseness of the pupil is normal.

D : Middle-nasal protrusion (2:02 - 4:00) - 1.96 %

D : Decentralization of the pupil is normal.

D : Ellipseness of the pupil is normal.

PARAMETERS OF THE AUTONOMIC NERVE WREATH (ANW):	S	D
Diameter (pix)	192	184
Perimeter (pix)	639	608
The ratio between Pupillary and Ciliary belts (%)	26.47	23.70
Normal (25..35%) or pathologic.	Normal	Spastic
Asymmetry of pupillary belt (normal: 0..5%)	8.33	1.09
	Pathology	Normal
Type of the ANW form -	regular	regular
	Normal	Normal

D: Frontal and basal zones of pupillary belt are constricted.

S: Lower nasal shift.

S: Frontal zone of pupillary belt is constricted.

Functional frustration in correlation between sympathetic and parasympathetic nervous systems. (The origin should be specified?)

Atonic colitis. Initial manifestations of pathology in compensation state.

Weakness of smooth muscles. (stable changes).

Increased emotional lability, predisposition to spastic reactions. Overstressed state.

PIGMENT SATURATED SUBTYPE.

For people with such subtype increased reactivity of lymphoid tissues (thymus, lymphatic nodes, tonsils, spleen, etc.) is typical.

Diseases of digestive tract mucosa, genitals are often observed.

Predisposition to immune system pathology (allergic diseases, collagenosis, reduced resistance to infections). Fast acetilator - sensitivity of the ferment systems to chemical and drug influences is increased.