



Personal Data

Name : HD_34
 Sex : male
 Age : 34

Numerical Data Analysis

PARAMETERS OF THE IRIS :	S	D
Diameter (pix) =	412	414
Area (pix)	133317	134614
PARAMETERS OF THE PUPILLARY BORDER :		
Diameter (pix)	100	112
Diameter of the pupil in relation to iris (%)	24	27
Normal for current age 23-32%	Normal	Normal
Pupil border deformation degree (normal:0%...5%)	5	2
PARAMETERS OF THE PUPIL RELATIVE TO THE IRIS :		
Distance between the pupil and iris centers (%)	7.28	6.76
Normal (lower than 5% of above) or pathology	Pathology	Pathology
PARAMETERS OF THE APPROXIMATE ELLIPSE		
Ellipseness degree of the pupil (normal: 95% ... 100%)	98	91
	Normal	Pathology
Pupil form type	chord	ellipse
	Pathology	Pathology
PARAMETERS OF THE PUPILLARY MARGIN :		
Type of the form -	regular	regular
	Normal	Normal

Diagnosis

S : Upper temporal protrusion (0:56 - 2:06) - 2.00 %
S : Lower temporal protrusion (4:04 - 5:36) - 4.00 %
S : Upper-nasal decentralization.
S : Ellipseness of the pupil is normal.

D : Middle-temporal flatness (7:50 - 10:12) - 16.07 %
D : Upper-nasal decentralization.
D : Oval-diagonal form of the pupil.

Tonsillo-cardial syndrome.
Liver disease.
Pancreas gland diseases.
Urogenital disturbances.

PARAMETERS OF THE AUTONOMIC NERVE WREATH (ANW):	S	D
Diameter (pix)	190	210
Perimeter (pix)	670	740
The ratio between Pupillary and Ciliary belts (%)	28.85	32.45
Normal (25..35%) or pathologic.	Normal	Normal
Asymmetry of pupillary belt (normal: 0..5%)	5.26	3.81
Type of the ANW form -	Pathology	Normal
	regular	lacerated
	Normal	Pathology

D: Frontal zone of pupillary belt is constricted.
S: Basal shift.
S: Frontal zone of pupillary belt is constricted.

Congestive symptoms in the small pelvis due to the chronic inflammatory diseases.
Weakness of smooth muscles. (stable changes).

Metabolic disturbance, decrease of elastotonic features.
Initial vertebral osteoarthrosis.

PIGMENT SATURATED SUBTYPE.

For people with such subtype increased reactivity of lymphoid tissues (thymus, lymphatic nodes, tonsils, spleen, etc.) is typical. Diseases of digestive tract mucosa, genitals are often observed. Predisposition to immune system pathology (allergic diseases, collagenosis, reduced resistance to infections). Fast acetilator - sensitivity of the ferment systems to chemical and drug influences is increased.