



Personal Data

Name : HK38
 Sex : female
 Age : 38

Numerical Data Analysis

PARAMETERS OF THE IRIS :	S	D
Diameter (pix) =	464	482
Area (pix)	169093	182467
PARAMETERS OF THE PUPILLARY BORDER :		
Diameter (pix)	102	108
Diameter of the pupil in relation to iris (%)	21	22
Normal for current age 23-32%	Miosis	Miosis
Pupil border deformation degree (normal:0%...5%)	0	3
PARAMETERS OF THE PUPIL RELATIVE TO THE IRIS :		
Distance between the pupil and iris centers (%)	8.19	8.30
Normal (lower than 5% of above) or pathology	Pathology	Pathology
PARAMETERS OF THE APPROXIMATE ELLIPSE		
Ellipseness degree of the pupil (normal: 95% ... 100%)	94	93
Pupil form type	Pathology ellipse	Pathology ellipse
	Pathology	Pathology
PARAMETERS OF THE PUPILLARY MARGIN :		
Type of the form -	regular	regular
	Normal	Normal

Diagnosis

S : Frontal flatness (11:36 - 1:14) - 3.92 %
S : Middle-temporal protrusion (2:02 - 3:56) - 3.92 %
S : Lower temporal protrusion (4:00 - 5:34) - 3.92 %
S : Middle-nasal protrusion (8:02 - 9:38) - 3.92 %
S : Upper-nasal decentralization.
S : Oval-horizontal form of the pupil.

D : Frontal flatness (10:50 - 1:04) - 3.70 %
D : Middle-nasal protrusion (2:08 - 4:42) - 5.56 %
D : Frontal decentralization.
D : Oval-horizontal form of the pupil.

Risk to blood cerebral hemisphere vessel spasm.
Pancreas gland disturbances.
Depression, blood cerebral circulation disturbance, Bronchial asthma
predisposition.

PARAMETERS OF THE AUTONOMIC NERVE WREATH (ANW):	S	D
Diameter (pix)	186	196
Perimeter (pix)	579	625
The ratio between Pupillary and Ciliary belts (%)	23.20	23.53
Normal (25..35%) or pathologic.	Spastic	Spastic
Asymmetry of pupillary belt (normal: 0..5%)	6.45	2.04
Type of the ANW form -	Pathology regular	Normal regular
	Normal	Normal

D: Frontal zone of pupillary belt is constricted.
S: Lower nasal shift.
S: Frontal zone of pupillary belt is constricted.

Parasympathotonic. Liminal sensitivity of nervous system is increased. Reflex activity is accelerated. Secretary and evacuation functions of digestive tract is heightened.

Increased emotional lability, predisposition to spastic reactions. Overstressed state.

PIGMENT SATURATED SUBTYPE.

For people with such subtype increased reactivity of lymphoid tissues (thymus, lymphatic nodes, tonsils, spleen, etc.) is typical. Diseases of digestive tract mucosa, genitals are often observed. Predisposition to immune system pathology (allergic diseases, collagenosis, reduced resistance to infections). Fast acetilator - sensitivity of the ferment systems to chemical and drug influences is increased.