



Personal Data

Name : MAY_40
 Sex : male
 Age : 40

Numerical Data Analysis

PARAMETERS OF THE IRIS :

	S	D
Diameter (pix) =	470	474
Area (pix)	173494	176460

PARAMETERS OF THE PUPILLARY BORDER :

Diameter (pix)	142	134
Diameter of the pupil in relation to iris (%)	30	28
Normal for current age 23-32%	Normal	Normal
Pupil border deformation degree (normal:0%...5%)	10	9

PARAMETERS OF THE PUPIL RELATIVE TO THE IRIS :

Distance between the pupil and iris centers (%)	4.26	6.33
Normal (lower than 5% of above) or pathology	Normal	Pathology

PARAMETERS OF THE APPROXIMATE ELLIPSE

Ellipseness degree of the pupil (normal: 95% ... 100%)	92	97
Pupil form type	Pathology ellipse	Normal circle
	Pathology	Normal

PARAMETERS OF THE PUPILLARY MARGIN :

Type of the form -	regular	regular
	Normal	Normal

Diagnosis

S : Frontal drawing in (11:10 - 1:26) - 1.41 %
S : Middle-temporal flatness (2:38 - 3:20) - 5.63 %
S : Lower temporal protrusion (3:24 - 5:24) - 7.04 %
S : Middle-nasal protrusion (7:46 - 9:04) - 4.23 %
S : Decentralization of the pupil is normal.
S : Oval-horizontal form of the pupil.

D : Middle-temporal drawing in (8:38 - 10:00) - 2.99 %
D : Upper-nasal decentralization.
D : Ellipseness of the pupil is normal.

Tonsillo-cardial syndrome.

Inguinal hernia(?)

Pathology of thyroid gland.

Liver disease.

Depression, blood cerebral circulation disturbance, Bronchial asthma predisposition.

PARAMETERS OF THE AUTONOMIC NERVE WREATH (ANW):	S	D
Diameter (pix)	244	226
Perimeter (pix)	755	743
The ratio between Pupillary and Ciliary belts (%)	31.10	27.06
Normal (25..35%) or pathologic.	Normal	Normal
Asymmetry of pupillary belt (normal: 0..5%)	3.28	3.54
	Normal	Normal
Type of the ANW form -	regular	regular
	Normal	Normal

D: Basal zone of pupillary belt is constricted.

S: Frontal zone of pupillary belt is constricted.

Weakness of smooth muscles. Initial manifestations of pathology in compensation state.

Chronic duodenitis. Secondary functional weakness of pancreas. Initial manifestations of pathology in compensation state.

Chronic colitis.

Increased emotional lability, predisposition to spastic reactions. Overstressed state.

Vertebral osteoarthritis.

Metabolism is disturbed.

HEMATOGENIC TETANIC SUBTYPE.

Increased reactivity of nervous system, predisposition to anxiety, fears, emotional stresses. Predisposition to thyrotoxicosis, migraine, cardialgia, neurasthenia, nettle rash, intestinal colic, impotence.

Neurosis, depressions, diseases of thyroid gland, bronchial asthma of neurogenic etiology, neurodermitis, eczema. Predisposition to ulcers of neurogenic origin. Dyskinesia of gallbladder is typical with the following development of cholelithiasis due to cholestasis.