



Personal Data

Name : MOA\_52\_  
 Sex : female  
 Age : 52

Numerical Data Analysis

	S	D
PARAMETERS OF THE IRIS :		
Diameter (pix) =	404	406
Area (pix)	128190	129462
PARAMETERS OF THE PUPILLARY BORDER :		
Diameter (pix)	78	80
Diameter of the pupil in relation to iris (%)	19	19
Normal for current age 21-25%	Miosis	Miosis
Pupil border deformation degree (normal:0%...5%)	34	26
PARAMETERS OF THE PUPIL RELATIVE TO THE IRIS :		
Distance between the pupil and iris centers (%)	10.40	5.42
Normal (lower than 5% of above) or pathology	Pathology	Pathology
PARAMETERS OF THE APPROXIMATE ELLIPSE		
Ellipseness degree of the pupil (normal: 95% ... 100%)	93	93
Pupil form type	Pathology ellipse	Pathology ellipse
	Pathology	Pathology
PARAMETERS OF THE PUPILLARY MARGIN :		
Type of the form -	regular	regular
	Normal	Normal

Diagnosis

S : Middle-temporal protrusion ( 2:30 - 3:54) - 2.56 %

S : Middle-nasal decentralization.

S : Oval-diagonal form of the pupil.

D : Middle-nasal protrusion ( 1:58 - 4:08) - 5.00 %

D : Upper-nasal decentralization.

D : Oval-horizontal form of the pupil.

Chronic inflammatory processes in nasopharynx.

Liver disease.

Risk to cardiospastic reaction.

Depression, blood cerebral circulation disturbance, Bronchial asthma predisposition.

Urogenital disturbances.

PARAMETERS OF THE AUTONOMIC NERVE WREATH (ANW):	S	D
Diameter (pix)	156	154
Perimeter (pix)	519	544
The ratio between Pupillary and Ciliary belts (%)	23.93	22.70
Normal (25..35%) or pathologic.	Spastic	Spastic
Asymmetry of pupillary belt (normal: 0..5%)	2.56	9.09
Type of the ANW form -	Normal drawn in Pathology	Pathology drawn in Pathology

D: Middle-temporal shift.

D: Frontal and basal zones of pupillary belt are constricted.

S: Frontal zone of pupillary belt is constricted.

Hypertension in the lesser circulation with the symptoms of cardio-vascular insufficiency.

Acquired colitis. Initial manifestations of pathology in compensation state.

Nephroptosis. (stable changes).

Changes in liver stroma. Decrease of detoxicative function.

Endogenic or exogenic intoxication.

Vertebral osteoarthrosis.

MIXED SUBTYPE WITH THE SIGNS OF URATE DIATHESIS.

Peculiar features of reactivity: normal function of lymphoid tissue (with some tendency to lowering), connective tissue weakness, predisposition to metabolic disturbances with deviation of acid-alkaline balance to the acid side.

Metabolic disturbances : osteochondrosis, arthrosis, lithogenesis.