



Personal Data

Name : NO26
 Sex : female
 Age : 26

Numerical Data Analysis

PARAMETERS OF THE IRIS :	S	D
Diameter (pix) =	420	420
Area (pix)	138544	138544
PARAMETERS OF THE PUPILLARY BORDER :		
Diameter (pix)	114	114
Diameter of the pupil in relation to iris (%)	27	27
Normal for current age 23-32%	Normal	Normal
Pupil border deformation degree (normal:0%...5%)	2	2
PARAMETERS OF THE PUPIL RELATIVE TO THE IRIS :		
Distance between the pupil and iris centers (%)	6.19	6.67
Normal (lower than 5% of above) or pathology	Pathology	Pathology
PARAMETERS OF THE APPROXIMATE ELLIPSE		
Ellipseness degree of the pupil (normal: 95% ... 100%)	100	98
	Normal	Normal
Pupil form type	circle	circle
	Normal	Normal
PARAMETERS OF THE PUPILLARY MARGIN :		
Type of the form -	regular	regular
	Normal	Normal

Diagnosis

S : Middle-nasal flatness (8:06 - 9:14) - 10.53 %
S : Upper-nasal decentralization.
S : Ellipseness of the pupil is normal.

D : Middle-nasal flatness (2:46 - 3:38) - 7.02 %
D : Frontal protrusion (10:56 - 0:06) - 1.75 %
D : Middle-temporal drawing in (8:04 - 9:10) - 1.75 %
D : Lower nasal protrusion (4:34 - 5:44) - 1.75 %
D : Upper-nasal decentralization.
D : Ellipseness of the pupil is normal.

Irritation of the ischiatic nerve.
Increased fatiguability.
Liver disturbances.
Pancreas gland diseases.

PARAMETERS OF THE AUTONOMIC NERVE WREATH (ANW):	S	D
Diameter (pix)	190	192
Perimeter (pix)	631	658
The ratio between Pupillary and Ciliary belts (%)	24.84	25.49
Normal (25..35%) or pathologic.	Spastic	Normal
Asymmetry of pupillary belt (normal: 0..5%)	1.05	8.33
	Normal	Pathology
Type of the ANW form -	regular	regular
	Normal	Normal

D: Middle-temporal shift.
D: Frontal and basal zones of pupillary belt are constricted.
S: Frontal and basal zones of pupillary belt are constricted.

Functional frustration in correlation between sympathetic and parasympathetic nervous systems. (The origin should be specified?)
Hypertension in the lesser circulation with the symptoms of cardio-vascular insufficiency.
Atonic colitis. (stable changes).

Changes in liver stroma. Decrease of detoxicative function.
Increased emotional lability, predisposition to spastic reactions. Overstressed state.
Initial vertebral osteoarthrosis.

PIGMENT SATURATED NEUROGENIC SUBTYPE.
People with such iris subtype have increased reactivity of nervous system and lymphoid tissue.
Diligence, purposefulness, high capacity for work is typical, which, however, may result in overstrain, nervous system exhaustion and nervous frustration. Migraine like headaches, irritation, fatigue, nervous and asthmatic attacks are frequent in such people, gastric and duodenum ulcer of nervous etiology is possible.