



Personal Data

Name : NT55
 Sex : male
 Age : 55

Numerical Data Analysis

| PARAMETERS OF THE IRIS : | S | D |
|--|--------------|-------------------|
| Diameter (pix) = | 398 | 418 |
| Area (pix) | 124410 | 137228 |
| | | |
| PARAMETERS OF THE PUPILLARY BORDER : | | |
| Diameter (pix) | 102 | 102 |
| Diameter of the pupil in relation to iris (%) | 25 | 24 |
| Normal for current age 21-25% | Normal | Normal |
| Pupil border deformation degree (normal:0%...5%) | 6 | 9 |
| | | |
| PARAMETERS OF THE PUPIL RELATIVE TO THE IRIS : | | |
| Distance between the pupil and iris centers (%) | 1.51 | 1.91 |
| Normal (lower than 5% of above) or pathology | Normal | Normal |
| | | |
| PARAMETERS OF THE APPROXIMATE ELLIPSE | | |
| Ellipseness degree of the pupil (normal: 95% ... 100%) | 98 | 92 |
| Pupil form type | Normal chord | Pathology ellipse |
| | Pathology | Pathology |
| | | |
| PARAMETERS OF THE PUPILLARY MARGIN : | | |
| Type of the form - | regular | regular |
| | Normal | Normal |

Diagnosis

S : Middle-nasal flatness (8:44 - 9:56) - 5.88 %
S : Middle-temporal flatness (2:22 - 3:42) - 13.73 %
S : Decentralization of the pupil is normal.
S : Ellipseness of the pupil is normal.

D : Lower nasal protrusion (2:34 - 7:52) - 1.96 %
D : Middle-temporal flatness (8:26 - 9:52) - 21.57 %
D : Decentralization of the pupil is normal.
D : Oval-vertical form of the pupil.

Chronic cardiac-lungs insufficiency.
Increased fatiguability.
Ischemic cardiac disease.
Circulatory cerebral disturbance with danger of ischemic variation.

| PARAMETERS OF THE AUTONOMIC NERVE WREATH (ANW): | S | D |
|---|-----------|---------|
| Diameter (pix) | 212 | 214 |
| Perimeter (pix) | 662 | 729 |
| The ratio between Pupillary and Ciliary belts (%) | 37.16 | 35.44 |
| Normal (25..35%) or pathologic. | Atonic | Atonic |
| Asymmetry of pupillary belt (normal: 0..5%) | 5.66 | 3.74 |
| | Pathology | Normal |
| Type of the ANW form - | regular | regular |
| | Normal | Normal |

S: Middle-temporal shift.
S: Basal zone of pupillary belt is constricted.
Sympathotonic. Liminal sensitivity of nervous system is decreased. Reflex activity is decelerated. Secretory and evacuation functions of digestive tract is reduced.
Overloads of the left ventricle.
Secondary functional weakness of pancreas. Initial manifestations of pathology in compensation state.

Chronic colitis.
Initial vertebral osteoarthrosis.

MIXED SUBTYPE WITH THE SIGNS OF URATE DIATHESIS.
Peculiar features of reactivity: normal function of lymphoid tissue (with some tendency to lowering), connective tissue weakness, predisposition to metabolic disturbances with deviation of acid-alkaline balance to the acid side.
Metabolic disturbances : osteochondrosis, arthrosis, lithogenesis.