


RESULTS OF THE IRIDOREFLEXOLOGIC EXAMINATION



○ Personal Data

Name : Deixy
 Sex : female
 Age : 55

Numerical Data Analysis

PARAMETERS OF THE IRIS :	S	D
Diameter (pix) =	448	430
Area (pix)	157633	145220
PARAMETERS OF THE PUPILLARY BORDER :		
Diameter (pix)	78	82
Diameter of the pupil in relation to iris (%)	17	19
Normal for current age 21-25%	Miosis	Miosis
Pupil border deformation degree (normal:0%...5%)	24	28
PARAMETERS OF THE PUPIL RELATIVE TO THE IRIS :		
Distance between the pupil and iris centers (%)	6.25	7.91
Normal (lower than 5% of above) or pathology	Pathology	Pathology
PARAMETERS OF THE APPROXIMATE ELLIPSE		
Ellipseness degree of the pupil (normal: 95% ... 100%)	97	93
Pupil form type	Normal chord	Pathology ellipse
	Pathology	Pathology
PARAMETERS OF THE PUPILLARY MARGIN :		
Type of the form -	regular	regular
	Normal	Normal

Diagnosis

S : Middle-nasal decentralization.
S : Ellipseness of the pupil is normal.

D : Basal protrusion (4:46 - 6:22) - 4.88 %
D : Upper-nasal decentralization.
D : Oval-vertical form of the pupil.

Insufficiency of valval apparatus of deep veins of lower extremities.
Liver disease.
Risk to cardiospastic reaction.
Circulatory cerebral disturbance with danger of ischemic variation.

PARAMETERS OF THE AUTONOMIC NERVE WREATH (ANW):	S	D
Diameter (pix)	166	178
Perimeter (pix)	531	570
The ratio between Pupillary and Ciliary belts (%)	23.78	27.59
Normal (25..35%) or pathologic.	Spastic	Normal
Asymmetry of pupillary belt (normal: 0..5%)	14.46	6.74
	Pathology	Pathology
Type of the ANW form -	regular	regular
	Normal	Normal

D: Lower temporal shift.
D: Frontal zone of pupillary belt is constricted.
S: Lower temporal shift.
S: Frontal zone of pupillary belt is constricted.

Functional frustration in correlation between sympathetic and parasympathetic nervous systems. (The origin should be specified?)
Hemodynamics disturbance in the vena cava inferior.
Nephroptosis. (stable changes).

Changes in liver stroma. Decrease of detoxicative function.
Initial vertebral osteoarthrosis.
Metabolism is disturbed. Intoxication.

PIGMENT SATURATED SUBTYPE.

For people with such subtype increased reactivity of lymphoid tissues (thymus, lymphatic nodes, tonsils, spleen, etc.) is typical.
Worsening of toxic agents withdrawal from the organism occurs and as a result congenital and acquired metabolic dysfunction and osteochondrosis.

ADVICE