


RESULTS OF THE IRIDOREFLEXOLOGIC EXAMINATION



Personal Data

Name : AD-17
 Sex : female
 Age : 17

Numerical Data Analysis

PARAMETERS OF THE IRIS :	S	D
Diameter (pix) =	430	428
Area (pix)	145220	143872
PARAMETERS OF THE PUPILLARY BORDER :		
Diameter (pix)	126	134
Diameter of the pupil in relation to iris (%)	29	31
Normal for current age 28-40%	Normal	Normal
Pupil border deformation degree (normal:0%...5%)	10	11
PARAMETERS OF THE PUPIL RELATIVE TO THE IRIS :		
Distance between the pupil and iris centers (%)	8.37	9.81
Normal (lower than 5% of above) or pathology	Pathology	Pathology
PARAMETERS OF THE APPROXIMATE ELLIPSE		
Ellipseness degree of the pupil (normal: 95% ... 100%)	95	97
	Normal	Normal
Pupil form type	chord	chord
	Pathology	Pathology
PARAMETERS OF THE PUPILLARY MARGIN :		
Type of the form -	regular	regular
	Normal	Normal

Diagnosis

S : Frontal flatness (11:10 - 1:22) - 7.94 %
S : Upper-nasal decentralization.
S : Ellipseness of the pupil is normal.

D : Lower nasal protrusion (3:36 - 5:26) - 4.48 %
D : Middle-temporal protrusion (8:48 - 9:50) - 1.49 %
D : Upper-nasal decentralization.
D : Ellipseness of the pupil is normal.

Absorbtion in large intestine is decreased.
Discirculatory encephalopathy. Depression.
Gallbladder insufficiency.
Inborn weakness of the pancreas.

PARAMETERS OF THE AUTONOMIC NERVE WREATH (ANW):	S	D
Diameter (pix)	216	228
Perimeter (pix)	781	752
The ratio between Pupillary and Ciliary belts (%)	29.61	31.97
Normal (25..35%) or pathologic.	Normal	Normal
Asymmetry of pupillary belt (normal: 0..5%)	6.48	4.39
	Pathology	Normal
Type of the ANW form -	regular	lacerated
	Normal	Pathology

D: Frontal and basal zones of pupillary belt are constricted.
S: Lower temporal shift.
S: Frontal and basal zones of pupillary belt are constricted.

Reduction of lungs ventilation.
Atonic colitis.

Increased emotional lability, predisposition to spastic reactions. Overstressed state. Initial vertebral osteoarthrosis.