



Personal Data

Name : JR-61
 Sex : female
 Age : 61

Numerical Data Analysis

PARAMETERS OF THE IRIS :	S	D
Diameter (pix) =	394	408
Area (pix)	121922	130741
PARAMETERS OF THE PUPILLARY BORDER :		
Diameter (pix)	94	102
Diameter of the pupil in relation to iris (%)	23	25
Normal for current age 21-25%	Normal	Normal
Pupil border deformation degree (normal:0%...5%)	9	3
PARAMETERS OF THE PUPIL RELATIVE TO THE IRIS :		
Distance between the pupil and iris centers (%)	8.12	5.39
Normal (lower than 5% of above) or pathology	Pathology	Pathology
PARAMETERS OF THE APPROXIMATE ELLIPSE		
Ellipseness degree of the pupil (normal: 95% ... 100%)	96	96
	Normal	Normal
Pupil form type	chord	circle
	Pathology	Normal
PARAMETERS OF THE PUPILLARY MARGIN :		
Type of the form -	regular	regular
	Normal	Normal

Diagnosis

S : Middle-temporal protrusion (2:38 - 4:24) - 4.26 %

S : Upper-nasal decentralization.

S : Ellipseness of the pupil is normal.

D : Frontal flatness (11:42 - 1:22) - 3.92 %

D : Middle-nasal decentralization.

D : Ellipseness of the pupil is normal.

Overloads of right parts of the heart.

Pulmonary pathology.

Pancreas gland disturbances.

PARAMETERS OF THE AUTONOMIC NERVE WREATH (ANW):	S	D
Diameter (pix)	174	176
Perimeter (pix)	691	640
The ratio between Pupillary and Ciliary belts (%)	26.67	24.18
Normal (25..35%) or pathologic.	Normal	Spastic
Asymmetry of pupillary belt (normal: 0..5%)	6.90	7.95
	Pathology	Pathology
Type of the ANW form -	regular	regular
	Normal	Normal

D: Lower temporal shift.

D: Frontal zone of pupillary belt is constricted.

S: Middle-temporal shift.

S: Frontal and basal zones of pupillary belt are constricted.

Functional frustration in correlation between sympathetic and parasympathetic nervous systems. (The origin should be specified?)

Overloads of the left ventricle.

Hemodynamics disturbance in the vena cava inferior.

Acquired colitis. Initial manifestations of pathology in compensation state.

Nephroptosis. (stable changes).

Acid-alkaline balance disturbance.

Initial vertebral osteoarthrosis.

Metabolism is disturbed.

LYMPHATIC SUBTYPE.

Increased reactivity of lymphoid tissues (thymus, lymphatic nodes, tonsils, spleen, etc).

Show the worsening of toxic agents withdrawal from the organism and, as a result, congenital and acquired metabolic dysfunctions.