



Personal Data

Name : DRVD-45
 Sex : male
 Age : 45

Numerical Data Analysis

PARAMETERS OF THE IRIS :	S	D
Diameter (pix) =	444	416
Area (pix)	154830	135918
PARAMETERS OF THE PUPILLARY BORDER :		
Diameter (pix)	86	72
Diameter of the pupil in relation to iris (%)	19	17
Normal for current age 23-32%	Miosis	Miosis
Pupil border deformation degree (normal:0%...5%)	12	42
PARAMETERS OF THE PUPIL RELATIVE TO THE IRIS :		
Distance between the pupil and iris centers (%)	4.05	5.77
Normal (lower than 5% of above) or pathology	Normal	Pathology
PARAMETERS OF THE APPROXIMATE ELLIPSE		
Ellipseness degree of the pupil (normal: 95% ... 100%)	97	92
Pupil form type	Normal	Pathology
	circle	ellipse
	Normal	Pathology
PARAMETERS OF THE PUPILLARY MARGIN :		
Type of the form -	regular	regular
	Normal	Normal

Diagnosis

S : Frontal drawing in (11:08 - 0:32) - 2.33 %
S : Middle-temporal protrusion (1:56 - 4:06) - 4.65 %
S : Basal protrusion (5:06 - 6:42) - 2.33 %
S : Decentralization of the pupil is normal.
S : Ellipseness of the pupil is normal.

D : Middle-nasal flatness (2:56 - 3:02) - 5.56 %
D : Upper-nasal decentralization.
D : Oval-diagonal form of the pupil.

Irritation of the ischiatic nerve.
Overloads of right parts of the heart.
Liver disturbances.
Urogenital disturbances.

PARAMETERS OF THE AUTONOMIC NERVE WREATH (ANW):	S	D
Diameter (pix)	166	152
Perimeter (pix)	543	520
The ratio between Pupillary and Ciliary belts (%)	22.35	23.26
Normal (25..35%) or pathologic.	Spastic	Spastic
Asymmetry of pupillary belt (normal: 0..5%)	2.41	9.21
	Normal	Pathology
Type of the ANW form -	regular	drawn in
	Normal	Pathology

D: Lower nasal shift
D: Frontal zone of pupillary belt is constricted.
S: Frontal zone of pupillary belt is constricted.

Parasympathotonic. Liminal sensitivity of nervous system is increased. Reflex activity is accelerated. Secretory and evacuation functions of digestive tract is heightened.
Weakness of smooth muscles. (stable changes).

Chronic colitis.
Predisposition to allergic reactions.
Metabolic disturbance, decrease of elastotonic features.
Initial vertebral osteoarthritis.

LYMPHATIC HYDROGENOUS SUBTYPE WITH SIGNS OF URATE DIATHESIS.
Violation of colloid balance in protein liquids, produced by the organism. Lymphoid tissue is highly reactive. Diseases, connected with metabolic disturbance are characteristic for the given subtype.
Predisposition to immunopathologic diseases (rheumatism, collagenosis) and metabolic disturbances (urolithic and cholelithic diseases, skin diseases).